FISH FACTS 2013

Results from fishery surveys conducted during the summer of 2013 are shown on the following pages. Fish capture techniques are dependent upon the objectives of the survey and may include gill nets or downhaul trap nets. Fish sizes are typically reported by weight (g) for gill nets and length (cm) for trap nets.

If you have any questions on survey methods or equipment, or if you have concerns about the health of a specific fishery, please contact your nearest Ministry of Environment fisheries biologist.

Waterbody and Location	Species Caught (Weight) and Population Status	Date Tested / Survey Gear / Remarks
Amisk Lake 54° 35' 102° 15' South of Creighton	Walleye: (48 - 3290 g) Fair Pike: (58 - 7190 g) Poor Perch: (35 - 163 g) Sparse Burbot: (160 - 2375 g)	July 15 - 18 / Gill nets. Large destination lake with use by residents, outfitters, campers, itinerant anglers, commercial fishery. Good angling reported. Majority of perch were forage size (≤18 cm).
Anglin Lake 55 - 27 W2 north of Prince Albert	Walleye: (135 - 1980 g) Moderate Pike: (275 - 1885 g) Fair Perch: (35 - 195 g) Poor	June 18 - 19 / Gill nets. No reports of angling success for walleye or pike at time of survey.
Berven Lake 55° 47' 104°33' via Hailstone Lake, Hwy 102 - north of La Ronge	Lake Trout: (80 - 3680 g) Sparse Pike: (340 - 2550 g) Fair Burbot: (590 - 1990 g)	July 30 - August 1 and August 6 - 8 / Gill nets. No reports of angling activity at time of survey.
Boire Lake 12,13,14 - 63 - 15 W3 northeast of Meadow Lake	Pike: (98 - 2092 g) Good Perch: (17 - 432 g) Moderate	August 19 - 20 / Gill nets. Pike - angling success reported as good, with larger size fish uncommon. Perch - winter angling success reported as good, with fair population of angling size fish (≥18 cm).
Boundary Reservoir 1 - 8 W2 south of Estevan	Walleye: (90 - 1060 g) Good Perch: (60 - 290 g) Poor Bullhead: (300 - 750 g)	July 29 - 31 / Gill nets. Walleye are more abundant than encountered in previous surveys. Capture of small walleye suggests recent successful reproduction. Pike populations remain sparse, as none were captured during the survey.
Christopher Lake 6, 7, 18 - 53 - 26 W2 north of Prince Albert	Walleye: (92 - 758 g) Good Pike: (388 - 1502 g) Moderate	June 19 - 20 / Gill nets. Angling success reported as fair at time of survey.
East Trout Lake 62 - 21, 22 W2 north of Candle Lake	Walleye: (565 - 2720 g) Poor Pike: (170 - 795 g) Sparse Lake Trout: (1485 - 2975 g) Poor Whitefish: (75 - 1740 g) Fair	August 27 - 29 / Gill nets. No angling reports available for lake trout, walleye or pike at time of survey.
Good Spirit Lake 29 - 30 - 5 W2 19 km southwest of Canora	Walleye: (120 - 1530 g) Good Pike: (700 - 5675 g) Poor Burbot: (2500 g)	June 25 - 26 / Gill nets. The walleye population appears to be less abundant than in the previous surveys, but is still good. Angling success was reported as excellent.
Granite Lake 54° 53' 102° 32' north of Hwy 106	Walleye: (60 - 2700 g) Fair Pike: (80 - 2240 g) Good Perch: (50 - 60 g) Sparse Burbot: (1940 - 2990 g)	June 3 - 5 / Gill nets. Campground and good boat launch present. Perch were forage size (≤18 cm).
Hailstone Lake 55° 48' 104° 33' 9.7 km north of Dickens Lake - Hwy 102	Pike: (760 - 1470 g) Moderate Perch: Sparse	June 25 - 26 / Gill nets. Angler use unknown, but believed to be light.

Herbert Reservoir	Walleye: (50 - 2400 g) Excellent	Sept 17 - 18 / Gill nets. The walleye
9 - 17 - 10 W3	Pike: (640 g) Sparse	population is very abundant, with most fish
5 km west of Herbert	Tike. (040 g) Sparse	weighing approximately 1 kg. Numerous
5 km west of Herbert		small walleye from the spring 2013 stocking
		were also caught.
Jumping Lake	Walleye: (390 - 1282 g) Good	July 9 - 10 / Gill nets. Stocked every second
44,45 - 24, 25 W2	walleye: (390 - 1282 g) Good	year with 250,000 walleye fry.
south of Birch Hills		year with 250,000 waneye fry.
	$\mathbf{P}_{\mathbf{r}} = \mathbf{F}_{\mathbf{r}} = \mathbf{f}_{\mathbf{r}} (242 - 276 + 1) \mathbf{F}_{\mathbf{r}}^{\dagger}$	Arra 15 16 / Cill rate Stanlad with 2 000
Junction Lake 36 - 66 - 25 W2	Brook Trout: (242 - 376 g) Fair Brown Trout: (118 - 762 g) Fair	Aug 15 - 16 / Gill nets. Stocked with 2,000 brook and 2,000 brown trout on an alternate
north of Weyakwin	Brown frout: (118 - 702 g) Fan	
norm of weyakwin		year bases. No reports of angling success at time of survey.
Lac Eauclaire	$\mathbf{L}_{\mathbf{a}} = \mathbf{T}_{\mathbf{a}} \mathbf{a} \mathbf{a} \mathbf{a} \mathbf{b} \mathbf{b} \mathbf{a} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} \mathbf{b} b$	June 19 - 20 / Gill nets. Lake Trout -
15,22,23 - 56 - 12 W3	Lake Trout: (1729 - 3651 g) Sparse Pike: (1164 - 3675 g) Poor	fingerling stocking discontinued. Pike - poor
north of Chitek Lake		population, however results are within the
north of Chitek Lake	Whitefish: (49 - 1904 g) Moderate	range of previous surveys. Angling success
	Perch: Forage size	for pike reported as fair. Whitefish are more
		abundant than encountered in previous
Loo Ilo o lo Cuerco	Wellower $(50, 2110, z)$ Second	surveys.
Lac Ile a la Crosse 55° 40' 107° 45'	Walleye: (50 - 2110 g) Sparse	July 3 - 6, July 8 - 11 / Gill nets. Walleye -
<i>33</i> 40 10/ ⁻ 45	Pike: (92 - 5070 g) Fair	angling reported as fair. Pike - angling
	Perch: (15 - 110 g)	reported as good. Perch were forage size $(<12, am)$
		(≤18 cm).
Lac La Peche	Walleye: (82 - 1216 g) Good	July 12 - 13 / Gill nets. No reports of
3, 10 - 46 - 8 W3	Pike: (112 - 1138 g) Good	angling success at time of survey.
west of Marcelin	Perch: (95 g) Sparse	O_{1} $(7, 9)$ (C')
Lake Lenore Town	Nil	Oct 7 - 8 / Gill nets. Survey conducted to
Reservoir 36 - 39 - 21 W2		determine rainbow trout stocking success.
		Nil catch suggests a sparse to non-existent
1 km west of Lake		trout population. Stocking to be temporarily
Lenore (Town)		discontinued if winter oxygen levels are
T	W-H (1260 - 2510 -) F	poor. Water level was low at time of survey.
Lepine Lake	Walleye: (1260 - 2510 g) Fair	August 14 - 15 / Gill nets. The walleye
28,29,30 - 63 - 24 W3 Meadow Lake	Pike: (130 - 1500 g) Good	population is more abundant than
	Perch : (40 - 50 g)	encountered in previous surveys with
Provincial Park		angling success reported as good. Pike - no
		larger size fish caught. Angling success for
		pike reported as good. Perch were forage $airse$ (<18 am)
Limestone Lake	Wellow: $(40, 1040, c)$ Cood	size (≤18 cm). June 17 - 19 / Gill nets. Popular destination
54° 38' 103° 13'	Walleye : (40 - 1040 g) Good	
	Pike : (210 - 2420 g) Moderate	with considerable angling activity - good
south of Hwy 106	Perch : (30 - 150 g) Sparse	success reported for walleye and pike. The
Little Leabfield Labo	Dought $(< 18 \text{ cm})$	majority of perch were forage size (≤ 18 cm). June 5 - 6 / Gill nets. Test netting conducted
Little Jackfish Lake	Perch: (\leq 18 cm)	
25,36 - 30 - 30 W1 Duck Mountain		to determine if perch population had
Duck Mountain Provincial Park		winterkilled. Prior to introduction of perch,
Lost Echo Lake	Wollows (51 2190 a) Madanata	the lake was managed for stocked trout. August 7 - 8 / Gill nets. Stocked with
26 - 59 - 19 W2	Walleye: $(51 - 2180 \text{ g})$ Moderate Pike: $(372 - 3175 \text{ g})$ Fair	200,000 walleye fry every second year. No
Narrow Hills Provincial	Pike: (372 - 3175 g) Fair Whitefish: (314 - 1914 g) Good	reports of angling success at time of survey.
Park	Whitefish: (314 - 1914 g) Good	reports of anging success at time of survey.
Madge Lake	Walleye: (90 - 1440 g) Moderate	July 8 - 10 / Gill nets. The pike population is
30, 31 - 30 W1	Pike: (210 - 1860 g) Good	more abundant than encountered in previous
Duck Mountain	Pike: $(210 - 1860 \text{ g}) \text{ Good}$ Perch: $(\leq 18 \text{ cm})$	surveys. Angling success for walleye
Provincial Park		reported as good. Perch were forage size
i i uvilicial f alk		(\leq 18 cm). Water level was extremely high at
		$(\leq 18 \text{ cm})$. Water level was extremely high at the time of the survey.
		the time of the survey.

Nasland Laba	N:1	Lung (7/Cill acts Test acting and ducted
Nesland Lake	Nil	June 6 - 7 / Gill nets. Test netting conducted
36 - 49 - 7 W3		to assess winterkill status of pike population.
west of Mont Nebo		To be stocked annually with 5,000 rainbow
		trout. Lake is aerated during the winter; be
		careful of thin ice.
Nickle Reservoir	Walleye: (50 - 1620 g) Good	Sept. 11 - 12 / Gill nets. An abundant
7 - 8 - 14 W2	Perch: (<100 - 730 g) Fair	population of walleye exists, with most fish
5 km southeast of		weighing approximately 750 g. Angling for
Weyburn		walleye has been good. Although no pike
		were caught during the survey, anglers report
		catching this species.
Picnic Lake	Perch: (<100 - 175 g)	June 3 - 4 / Gill nets. Test netting conducted
11 - 49 - 19 W3		to determine if perch population had
6 km east of Edam		winterkilled. Majority of perch were forage
		size (≤ 18 cm).
Pierce Lake	Walleye: (365 - 2260 g) Poor	August 12 - 14 / Gill nets. The walleye, pike
63,64 - 25 W3	Pike: (210 - 6890 g) Fair	and perch populations are more abundant
Meadow Lake	,	than encountered in previous surveys.
Provincial Park	Lake Trout: (805 g) Sparse	
FIOVINCIAI PATK	Perch: (35 - 115 g) Fair	Angling success for walleye and pike
		reported as good. Lake trout – population is
		similar to previous surveys. Angling success
		reported as good. The majority of perch
		were forage size (≤18 cm).
Pine Lake	Splake: (878 - 1170 g) Poor	July 2 - 3 / Gill nets. Stocked every second
18 - 58 - 18 W2		year with 4,000 splake.
Narrow Hills Provincial		
Park		
Piprell Lake	Splake: (352 - 940 g) Poor	June 20 - 21 / Gill nets. Fair to good angling
59 - 20 W2	Brown Trout: (150 - 814 g) Poor	success for all trout species reported at time
north of Whiteswan	Rainbow Trout: (214 - 412 g) Moderate	of survey.
Lake	Tiger Trout: (264 - 644 g) Poor	
Rafferty Reservoir	Walleye: (70 - 2510 g) Good	Aug 19 - 22 / Gill nets. Good range in sizes
2 - 9 W2	Pike: (450 - 2420 g) Fair	of walleye. Number of walleye caught is
northwest of Estevan	Perch: (30 - 480 g) Moderate	similar to previous surveys. Majority of
northwest of Lste van	rerent (30 - 400 g) Woderate	perch caught were angling size (≥ 18 cm).
	NI:1	
Ridge Lake	Nil	July 4 - 5 / Gill nets. Stocked every second
11 - 61 - 21 W2		year with 4,000 brook trout.
northwest of Piprell		
Lake		
Scythes Lake	Walleye: (157 - 1795 g) Sparse	May 27 - 29 / Gill nets. No reports of
55° 24' 104° 55'	Pike : (157 - 995 g) Fair	angling success at time of survey. The
via Little Deer Lake -	Perch : (60 - 121 g) Sparse	majority of perch were forage size (≤ 18 cm).
Hwy 102, north of La		
Ronge		
Shell Lake	Pike: (90 - 2190 g) Good	June 4 - 5 / Gill nets. Pike are more abundant
5,8,17 - 56 - 11 W3		than encountered in previous surveys, with
northeast of Chitek Lake		angling success reported as fair. Larger size
		pike are uncommon. Perch - although
		angling size perch have been present in
		previous surveys, none were caught in this
G 100 1 2 1		previous surveys, none were caught in this survey - one was found in a pike stomach.
Smallfish Lake	Pike: (118 - 292 g) Fair	previous surveys, none were caught in this survey - one was found in a pike stomach.July 23 - 24 / Gill nets. No reports of angling
39,40 - 30 W1	Pike: (118 - 292 g) Fair Perch: (90 - 292 g) Fair	previous surveys, none were caught in this survey - one was found in a pike stomach.
		previous surveys, none were caught in this survey - one was found in a pike stomach.July 23 - 24 / Gill nets. No reports of angling
39,40 - 30 W1 southeast of Hudson Bay	Perch: (90 - 292 g) Fair	previous surveys, none were caught in this survey - one was found in a pike stomach. July 23 - 24 / Gill nets. No reports of angling success at time of survey.
39,40 - 30 W1 southeast of Hudson Bay Spirit Lake	Perch: (90 - 292 g) Fair Walleye: (304 - 2206 g) Good	 previous surveys, none were caught in this survey - one was found in a pike stomach. July 23 - 24 / Gill nets. No reports of angling success at time of survey. July 25 - 25 / Gill nets. Anglers report fair to
39,40 - 30 W1 southeast of Hudson Bay	Perch: (90 - 292 g) Fair	previous surveys, none were caught in this survey - one was found in a pike stomach. July 23 - 24 / Gill nets. No reports of angling success at time of survey.

Struthers Lake 2, 3 - 45 - 23 W2 southeast of Birch Hills Sturgeon Lake 51 - 1, 2 W3	Walleye: (583 - 794 g) Poor Pike: (1064 g) Poor Perch: (39 - 48 g) Sparse Walleye: (138 - 1830 g) Moderate Pike: (608 - 1348 g) Poor	July 30 - 31 / Gill nets. Stocked every second year with 200,000 walleye fry. June 27 - 28 / Gill nets. Stocked every second year with 1.4 million walleye fry.
north of Holbien	Perch: (80 - 222 g) Sparse	second year what i. I minion waneye ny.
Swede Lake 14 - 57 - 6 W3 northeast of Big River	Pike: (187 - 3070 g) Moderate	June 18 - 19 / Gill nets. Pike - population has increased compared to previous surveys. Angling success unknown. No perch were caught; however forage size fish were observed at boat launch.
Swift Current Reservoir 18 - 15 - 13 W3 1 km south of Swift Current	Walleye: (490 - 1040 g) Fair Pike: (4360 g) Poor Perch: (150 - 250 g) Moderate	Sept 18 - 19 / Gill nets. Despite its small size and shallowness, this water body contains numerous species of game fish. All perch caught were angling size (≥18 cm).
Tyrell Lake 54° 54' 102° 07' north of Hwy 106	Pike : (92 - 7598 g) Good Perch : (31 - 477 g) Fair	June 5 - 6 / Gill nets. Extensive shallow area at boat ramp makes launching difficult.
White Bear Reservoir 24 - 22 - 17 W3 1 km northwest of White Bear (Hamlet)	Rainbow Trout: (400 - 750 g) Moderate	Sept 16 - 17 / Gill nets. Test netting conducted to check survival of trout and to determine if stocking should continue. Evidence of angler activity at time of survey.