Fish Facts 2022



The results from the 2022 fishery surveys are available in the chart below. Lakes are mainly surveyed using two types of equipment; gill nets or downhaul trap nets. Both types of equipment are effective in capturing fish, however, fish caught in downhaul traps can generally be released after being measured, whereas fish caught in gill nets are generally not releasable. For this reason, the size of fish caught in downhaul trap nets is recorded by total length to minimize stress and harm to the fish, while size of fish captured in gill nets is reported by weight.

If you have any questions on survey methods or equipment, or if you have concerns about the health of a specific fishery, please contact the ministry's Inquiry Centre at 1-800-567-4224.

Waterbody and Location	Species Caught (weight or length) and Population Status	Date Tested/Survey Gear/Remarks
Broderick Reservoir	Walleye: (105 - 3720 g) Sparse	June 6 - 7. Gill nets. Consistent walleye
51° 28′ 106° 56′	Pike: (185 - 2760 g) Good	abundance, with an increase in pike and
south of Broderick	Perch: (30 - 130 g) Good	perch relative to previous survey. Pike
	Whitefish: (1200 - 2720 g) Good	population is sustained by natural
		reproduction. Most perch were of
		forage size (<18.0 cm).
Diefenbaker Lake	Walleye: (48 - 4800 g)	July 18 - 27. Gill nets. Increase in
50° 42′ 107° 26′	Sauger: (59 - 1035 g)	walleye, sauger, goldeye and lake
south of Saskatoon	Pike: (50 - 5920 g)	sturgeon abundance relative to the
	Perch: (30 - 450 g)	previous 2011 survey, with walleye the
	Whitefish: (84 - 2575 g)	second most common species caught.
	Goldeye: (55 - 1410 g)	Whitefish, pike, burbot and cisco
	Burbot: (220 – 2650 g)	abundance is comparable to the
		previous survey, with whitefish the most
		common fish species caught. Perch
		continue to decline. No trout of any
		species were caught during the survey.
Fishing Lake	Walleye: (80 - 2162 g) Excellent	June 21 - 23. Gill nets. Historic walleye
(Big and Little)	Pike: (680 - 2245 g) Fair	abundance with natural reproduction
51° 50′ 103° 32′	Perch: nil catch	occurring and 18-year classes present.
south of Kuroki		Slight decrease in pike abundance
		compared to previous survey with
		natural reproduction occurring and
		eight-year classes present.
Lac la Ronge	Lake trout: (360 - 5350 g)	Aug 22 - 26. Gill nets. SPIN (Summer
55° 12′ 105° 03′	Burbot: (370 - 3650 g)	Profundal Index Netting) assessment.
Nut Bay, Bear Island		Lake trout population estimate and size
		range is similar to initial SPIN survey in
		2016.

^{*}SPIN netting is a rapid assessment gill netting method that is specifically used to collect information about lake trout.

Madaa Laka	Welleyer (CO 2410 a) Fair	Lune 15 17 Cill note Decrees in
Madge Lake	Walleye: (60 - 2410 g) Fair	June 15 - 17. Gill nets. Decrease in
51° 40′ 101° 38′	Pike: (386 - 7550 g) Good	walleye abundance from previous
Duck Mountain	Perch: (35 - 105 g) Poor	surveys with 16-year classes present.
Provincial Park		Natural reproduction occurring with
		strong three-year-old age class linked to
		2019 fry stocking. Increase in pike
		abundance from previous survey with
		10-year classes present. Natural
		reproduction occurring. Decrease in
		perch abundance, with most perch of
		forage size (<18.0 cm).
Peck Lake	Walleye: (33 - 2340 g) Moderate	August 2 - 5. Gill nets. Walleye
53° 54′ 109° 36′	Pike: (500 - 4480 g) Poor	abundance has slightly increased since
northwest of Turtle	Perch: (26 - 134 g) Fair	the previous survey and is the highest
Lake	Whitefish: (150 - 1850 g) Poor	on record. Whitefish abundance has
	Burbot: (70 - 1740 g) Sparse	improved since the previous survey but
		remains well below the historical range.
		Pike abundance has decreased since the
		previous survey.
Round Lake (Nisbet)	Perch: (30 - 140 g) Poor	June 16 - 17. Gill nets. The lake is subject
53° 20′ 106° 01′	Pike: (1800 – 2750 g) Poor	to occasional winterkill. Survey results
northwest of Prince		indicate that the lake likely experienced
Albert.		a partial winterkill in 2021.
Waterhen Lake	Walleye: (134 - 3050 g) Fair	August 8 - 12. Gill nets. Walleye and pike
54° 28′ 108° 25′	Pike: (166 - 2800 g) Fair	abundance decreased compared to the
east of Cold Lake	Perch: (42 - 130 g) Fair	previous survey, but both remain within
	Whitefish: (350 - 1780 g) Fair	their historical ranges. Perch abundance
	Burbot: (600 - 1480 g) Sparse	remains high relative to their historical
		range across all previous surveys.